

## STATUS OF INSECTIVOROUS PLANTS IN NORTHEAST INDIA

PRAVEEN KUMAR VERMA • Shifting Cultivation Division • Rain Forest Research Institute • Sotai Ali • Deovan • Post Box # 136 • Jorhat 785 001 (Assam) • India • pkverma\_bryo@yahoo.co.in  
JAN SCHLAUER • Zwischenstr. 11 • 60594 Frankfurt/Main • Germany • jan@carnivorousplants.org  
KRISHNA KUMAR RAWAT • CSIR-National Botanical Research Institute • Rana Pratap Marg • Lucknow -226 001 (U.P) • India  
KRISHNA GIRI • Shifting Cultivation Division • Rain Forest Research Institute • Sotai Ali • Deovan • Post Box #136 • Jorhat 785 001 (Assam) • India

Keywords: Biogeography, India, diversity, Red List data.

### Introduction

There are approximately 700 identified species of carnivorous plants placed in 15 genera of nine families of dicotyledonous plants (Albert *et al.* 1992; Ellison & Gotelli 2001; Fleischmann 2012; Rice 2006) (Table 1).

In India, a total of five genera of carnivorous plants are reported with 44 species; *viz.* *Utricularia* (38 species), *Drosera* (3), *Nepenthes* (1), *Pinguicula* (1), and *Aldrovanda* (1) (Santapau & Henry 1976; Anonymous 1988; Singh & Sanjappa 2011; Zaman *et al.* 2011; Kamble *et al.* 2012). Interestingly, northeastern India is the home of all five insectivorous genera, namely *Nepenthes* (commonly known as tropical pitcher plant), *Drosera* (sundew), *Utricularia* (bladderwort), *Aldrovanda* (waterwheel plant), and *Pinguicula* (butterwort) with a total of 21 species. The area also hosts the “ancestral false carnivorous” plant *Plumbago zelayanica*, often known as murderous plant.

### Climate

Lowland to mid-altitude areas are characterized by subtropical climate (Table 2) with maximum temperatures and maximum precipitation (monsoon) in summer, *i.e.*, May to September (in some places the highest temperatures are reached already in April), and average temperatures usually not dropping below 0°C in winter. As usual average temperatures decrease with increasing elevation, and in highlands, a mountain/alpine climate prevails. In Sikkim and Arunachal Pradesh, the tree line ranges from 5500 m in the north to 4000 m in the south, and the snow line ranges from 6100 m in the north to 4900 m in the south.

### Chorology

The extreme northeast of India (Sikkim, Darjeeling – the northern part of West Bengal, Arunachal Pradesh, Assam, Meghalaya, Nagaland, Manipur, Tripura, Mizoram) is one of the most floristically most diverse places in Asia as it is located in the contact zone between the temperate flora from high elevations (Himalaya), the tropical flora from lowland peninsular India and Burma, and the subtropical flora from Yunnan, China. To the west it shares a border with Nepal, to the north with Bhutan, and China (Tibet), to the east with Burma (Myanmar) and to the south with Bangladesh. Covering less than 275,000 km<sup>2</sup>, it is composed of six different floristic provinces in four

Table 1. Comparison of carnivorous plant diversity (world/India/NE India). Data derived from the Carnivorous Plant Database. [http://www.omnisterra.com/bot/cp\\_home.cgi](http://www.omnisterra.com/bot/cp_home.cgi), accessed 24 Feb. 2014.

Order	Family	Genus	Number of described species		
			World	India	NE India
Oxalidales	Cephalotaceae	<i>Cephalotus</i>	1		
Nepenthales (Caryophyllales s. lat.)	Droseraceae	<i>Drosera</i>	168	3	2
		<i>Aldrovanda</i>	1	1	1
		<i>Dionaea</i>	1		
	Drosophyllaceae	<i>Drosophyllum</i>	1		
	Nepenthaceae	<i>Nepenthes</i>	127	1	1
	Dioncophyllaceae	<i>Triphyophyllum</i>	1		
Ericales	Sarraceniaceae	<i>Darlingtonia</i>	1		
		<i>Heliamphora</i>	20		
		<i>Sarracenia</i>	8		
	Roridulaceae	<i>Roridula</i>	2		
Scrophulariales (Lamiales)	Byblidaceae	<i>Byblis</i>	7		
	Lentibulariaceae	<i>Pinguicula</i>	94	1	1
		<i>Genlisea</i>	27		
		<i>Utricularia</i>	234	38	16
Total			693	44	21

Regions according to Takhtajan's (1986) scheme: the Northern Burmese, Eastern Himalayan, and Khasi-Maipur Provinces belonging to the Eastern Asiatic Region, the Tibetan Province belonging to the Irano-Turanian Region, the Bengal Province belonging to the Indian Region, and the Southern Burmese Province belonging to the Indochinese Region. Any area of comparable size and latitudinal extension (between 22°N and 30°N) in the New World or in Africa would cover at the most three Provinces in at the most two Regions. This extraordinary diversity is mainly due to the unique topographical profile, ranging from almost sea level in the Brahmaputra valley to 8598 m elevation (Kangchenjunga) at the Nepal border.

*Pinguicula alpina* (arctic + alpine + Himalayan) belongs to the temperate element. The paleotropical element is represented by *Nepenthes khasiana*. Its entire range is contained in the area of concern and is located north of the Tropic of Cancer, *i.e.*, this is the only species in the genus *Nepenthes* naturally occurring outside the tropics. *Drosera peltata* is widespread in eastern Asia, Australia, and Oceania (Gibson *et al.* 2012, as *D. lunata*). It is the only tuber forming sundew to reach beyond the Australian diversification center of this group. *Drosera burmannii* is widespread in the Australasian tropics. While it is disputed whether its only record from Africa (Sierra Leone) is native there, another, closely related species (*D. sessilifolia*) is its neotropical counterpart. The tropical bladderworts in the area are the terrestrial *Utricularia hirta*, *U. pubescens*, *U. caerulea*, *U. bifida*, *U. recta* (syn. *U. scandens* subsp. *firmula*), *U. foveolata*, *U. subulata*, and the aquatic *U. aurea*, *U. stellaris*, and *U. inflexa*, all of which are more or less widespread in tropical eastern Asia

Table 2. Climate data of representative towns in NE India (classification according to Köppen & Geiger 1930-1939, data from Wikipedia <a href="http://en.wikipedia.org/">http://en.wikipedia.org/</a> ).			
Location; State; Elevation	Climate Class	Precipitation	Temperature
		Annual /Monthly	Summer / Winter
Aizawl; Mizoram; 1132 m	Humid Subtropical (Cwa)	2200 mm / 6 mm (Jan.) - 305 mm (Aug.)	18-27°C (max.: Apr.) / 25-11°C
Agartala; Tripura; 13 m		2200 mm / 8 mm (Dec.) - 455 mm (Jun.)	24- 34°C (max.: Apr.) / 33-10°C
Guwahati; Assam; 55 m		1700 mm / 7 mm (Dec.) - 345 mm (Jul.)	22-32°C / 30-10°C
Imphal; Manipur; 790 m		1400 mm / 12 mm (Jan.) - 225 mm (Jun.)	15-33°C / 30-11°C
Kohima; Nagaland; 1450 m		1800 mm / 8 mm (Dec.) - 370 mm (Jul.)	17-25°C / 23-5°C
Shillong; Meghalaya; 1500 m	Subtropical- Highland (Cwb)	2200 mm / 9 mm (Dec.) - 470 mm (Jun.)	15-24°C / 23-4°C
Itanagar; Arunachal Pradesh; 750 m		2300 mm / 15 mm (Nov.) - 510 mm (Jul.)	18-34°C / 18-8°C
Darjeeling; N West Bengal; 2050 m		3000 mm / 7 mm (Dec.) - 780 mm (Jul.)	11-20°C / 14-2°C
Gangtok; Sikkim; 1700 m		3600 mm / 23 mm (Dec.) - 650 mm (Jul.)	13-22°C / 18-4°C

and many spread to tropical Africa and Australia, while only *U. pubescens* and *U. subulata* are also found in the Americas.

*Utricularia gibba* and *Aldrovanda vesiculosa* represent azonal aquatics not closely tied to any climatic vegetation zone.

An exceptionally large number of species of the subtropical to tropical section *Phyllaria* of *Utricularia* (*U. furcellata*, *U. christopheri*, *U. brachiata*, *U. striatula*, *U. multicaulis*) are native or even endemic to this area. *Utricularia kumaonensis*, *U. salwinensis*, and *U. forrestii*, likewise members of this section, are known from the Salween-Irrawaddy divide in westernmost Yunnan, China/northernmost Burma, *i.e.*, within some 200 km to the east of the area treated here. As all three species do also occur in the immediate vicinity to the north (Nepal, Bhutan, Tibet), it is quite probable that they occur in Himalayan northeastern India although no specimens have apparently been collected from here so far.

Other species presently not recorded from the area but known from surrounding territories are: *Drosera indica* (paleotropical, including, West Bengal, *cf.* Basak 1975, China [Yunnan], Burma); *Utricularia capillacea* (syn. *U. scandens*, paleotropical, including, Bihar, Nepal, China [Tibet, Yunnan], Burma, Bangladesh); *U. graminifolia* (tropical East Asian, including, China [Yunnan], Burma); *U. uliginosa* (tropical East Asian, including, China [Yunnan], Burma); *U. australis* (azonal, widespread in the Old World, including, Nepal, Bhutan, China [Tibet, Yunnan], Burma); *U. minor* (circumboreal, including, Nepal, Bhutan, China [Tibet, Yunnan], Burma).

The majority (some 20) of the remaining Indian bladderwort species not known from the extreme northeast are members of *U.* section *Oligocista* and are more or less restricted to the Deccan peninsula, predominantly in the Western Ghats (Janarthanam & Henry 1992; Taylor 1989).

#### Conservation Status

**1. *Nepenthes khasiana* Hook. f.:** Popularly known as tropical pitcher plants or monkey cups, *Nepenthes khasiana* belongs to the monotypic family Nepenthaceae with 127 natural and hybrid species across the world (Jebb & Cheek 1997; Wikipedia 2013). It is the only native Indian species of insectivorous plant which is an endemic of Garo, Khasi, and Jaintia hills of Meghalaya. It greatly varies in its habit ranging from prostrate to scandent or rarely erect herbs, undershrubs, or shrubs. The plants are mostly associated with the natural vegetation of very humid climates.

**Distribution in northeast India:** Endemic to Meghalaya: East Khasi hills, West Khasi hills, Jaintia hills, south Garo hills, west Garo hills (Joseph & Joseph 1986; Venugopal & Devi 2003; Singh *et al.* 2011).

**Present status:** Habitat destruction is one of the main threats especially in Jaintia and Garo hills, where the coal mining plays a significance role in the decline of its natural population (Prasad & Jeeva 2009; Singh *et al.* 2011). Excessive collection, developmental activities, deforestation, fragmentation, increasing production of waste and pollutants, forest fire, *jhum* cultivation, and poor seed germination ability of the plant are also some of the causes of declining populations (Bordoloi 1977; Jain & Sastry 1980; Rathore *et al.* 1991; Mandal & Mukherjee 2011). It is classified as 'Endangered' (EN) and is incorporated in the Appendix I of CITES and Negative List of Exports of the Government of India (Ziemer 2010; Mandal & Mukherjee 2011). The species is restricted only in some pockets of Meghalaya.

**2. *Drosera* L.:** One of the largest genera of carnivorous plants (commonly known as sundews) with more than 160 species belonging to the family Droseraceae (Rice 2006; Jayaram & Prasad 2006).

**2.1. *Drosera burmannii* Vahl:** One of the most common insectivorous species among Indian carnivorous plants and found in tropical and subtropical parts of the country. This annual species possess spatulate leaves and forms rosettes which spans up to 2.5 cm. This species has important medicinal properties as it contains several medicinally active compounds including quinones (plumbagin), hydroplumbagin glucoside, flavonoids (kaempferol, myricetin, quercetin, and hyperoside), rossolidside (7-methyl-hydrojuglone-4-glucoside), and has been used for several disorders (Raju & Christina 2013).

**Distribution:** Arunachal Pradesh, Assam, Meghalaya, Sikkim, Tripura (Majumdar *et al.* 2011).

**Present status:** Least concerned (LC) ver.3.1, IUCN, 2011 (Zhuang 2011a). The species has a wide range of distribution and no current threats are reported for Indian populations.

**2.2 *Drosera peltata* Thunb. (syn. *Drosera lunata* Buch.-Ham. ex DC.):** This is a climbing or scrambling sundew species commonly called shield sundew. The species possesses cauline or peltate cauline leaves and can be easily identified in the field with its aerial parts growing from 5 to 15 cm in height. The species is a promising medicinal plant species due to its antimicrobial activity particularly against the oral bacteria (Didry *et al.* 1998).

**Distribution in northeast India:** Meghalaya (Jaintia hills, east and west Khasi hills), (Venugopal *et al.* 2007).

**Present status:** The species is somewhat rare and known only in some areas of Meghalaya growing on exposed rocks in association with mosses.

**3. *Utricularia* L.:** Another large genus of carnivorous plants with more than 230 species, which belongs to the family Lentibulariaceae (Parnell 2005; Fleischmann 2012) and commonly grows in floating as well as marshy lands. Unlike the other carnivorous plants discussed here, *Utricularia* often lives in aquatic conditions. A total of 16 species are found in northeastern India.

**3.1. *Utricularia aurea* Lour.:** Commonly known as golden bladderwort and is common in marshy and wet land of the area. Another important plant with compounds showing anti-tumor activity have been reported (Choosawad *et al.* 2005).

**Distribution in northeast India:** Assam (Sharma 2012, as *Utricularia flexuosa* Vahl.), Tripura (Bhowmik & Datta 2013).

**Present status:** The IUCN, 2103 listed the species under Least Concern (LC) ver 3.1 (IUCN 2013).

**3.2. *Utricularia bifida* L.:** The small and annual plants are found growing in marshy land as well as damp soil and commonly known as bifid bladderwort.

**Distribution in northeast India:** Assam, Meghalaya, Tripura (Bhowmik & Datta 2012a, 2013).

**Present status:** No major threats has been reported regionally as well as globally hence it been identified and is therefore listed as Least Concern (LC) 3.1. (Zhuang 2011b).

**3.3. *Utricularia brachiata* Oliv.:** Discovered by J.D. Hooker from Sikkim Himalaya and partially described by D. Oliver in 1859. The plants are very small, inconspicuous, with delicate, reniform leaves and grow as epiphyte intermixed with bryophytes on tree trunks (Compton 1909).

**Distribution in northeast India:** Arunachal Pradesh, Sikkim- Kyongnosla Alpine Sanctuary (Compton 1909; Taylor 1989).

**Present status:** In northeast India the species is very rare and restricted only to the upper hills of eastern Himalaya with in rather small populations.

**3.4. *Utricularia caerulea* L.:** Small to medium sized species commonly known as blue bladderwort (Kato *et al.* 2006; Kamble *et al.* 2012).

**Distribution in northeast India:** Meghalaya, Tripura (Bhowmik & Datta 2012b).

**Present status:** Very common in distribution and found from Madagascar to Japan and Australia through India and Malaysia (Kamble *et al.* 2012).

**3.5. *Utricularia christopheri* P.Taylor:** Perennial in nature.

**Distribution in northeast India:** Confined to East Himalayan region including Nepal; Sikkim- Changu, Lachen (Taylor 1989).

**Present status:** Grows in very high elevation in Sikkim Himalaya and Nepal with very limited distribution (Taylor 1989).

**3.6. *Utricularia gibba* L.:** A perennial aquatic herb commonly known as humped bladderwort. May be floating, growing submerged, or exposed on wet soil. One of the most common species among *Utricularia* (Taylor 1989).

**Distribution in northeast India:** Meghalaya (as *U. khasiana* Joseph & Mani), Tripura (as *U. gibba* subsp. *exoleta* (R. Br.) P.Taylor) (Deb 1983; Bhowmik & Datta 2013), Assam and Manipur (as *U.*

*exoleta* R. Br.) (Dixit & Bera 2012; Hazarika & Borthakur 2012).

**Present status:** Least Concern (LC) ver 3.1 (Anitha 2011).

**3.7. *Utricularia foveolata* Edgew.:** A small short-lived herb growing in shallow water or wet soil in seasonally wet grassland or occasionally as a weed in ricefields.

**Distribution in northeast India:** Meghalaya: Elephant Falls as *U. baouleensis* A. Chev. (Subramanyam & Kamble 1968).

**Present status:** Very rare in occurrence as it has been reported only once from Elephant Falls of Meghalaya in 1968 and no further reports are available from this area. Hence more detailed surveys are necessary to reveal its current status in India.

**3.8. *Utricularia furcellata* Oliv.:** Grows on moist logs or rocks along with mosses and form 3-5 mm wide rosettes.

**Distribution in northeast India:** Assam, Meghalaya, Mizoram, West Bengal (Suksathan & Parnell 2010; Chew & Haron 2011; Panday *et al.* 2013; Yee *et al.* 2012).

**Present status:** The status is yet to be worked out; however, recently it has been considered as rare in Mizoram (Panday *et al.* 2013).

**3.9. *Utricularia hirta* J.G.Klein ex Link.:** A small perennial, terrestrial insectivorous plant, native of Southeast Asia, and always growing in damp open places.

**Distribution in northeast India:** Meghalaya (Joseph & Mani 1982, as *U. tayloriana* J. Joseph & J. Mani), Tripura (Joseph & Mani 1982; Natarajan *et al.* 2008).

**Present status:** Least Concern (LC) ver 3.1 (Anitha 2011).

**3.10. *Utricularia multicaulis* Oliv.:** Very small growing annually and native of Southeast Asia.

**Distribution in northeast India:** Sikkim (Oliver 1859).

**Present status:** Discovered from Sikkim by Oliver (1859). Though the species has also been reported from China, in India, the plants seem extremely rare, as further reports are scanty.

**3.11. *Utricularia striatula* Sm.:** Commonly known as striped bladderwort is a small insectivorous herb which is growing on wet rock or even on tree trunks.

**Distribution in northeast India:** Meghalaya, Nagaland, Sikkim (Lal *et al.* 2003; Lansdown *et al.* 2013).

**Present status:** Least Concern (LC) ver 3.1.

**3.12. *Utricularia pubescens* Sm.:** A small to medium sized herbaceous annual plant growing on soil as well as on wet rocks.

**Distribution in northeast India:** Meghalaya (Rao & Joseph 1967).

**Present status:** Reported only once from the area (*v.s.*). Seems to be rare, as it is known from only one more locality in India, *i.e.* Rajpur, near Dehradun (Saxena 1965).

**3.13. *Utricularia recta* P. Taylor (*U. scandens* subsp. *firmula* (Oliver) Z. Yu Li):** Small herb, growing in grassland as well as on rocks.

**Distribution in northeast India:** NE India (Li & Cheek 2011).

**Present status:** Widespread in Africa and Asia. In India in the Himalayan part of Uttarakhand (Kumaun), in Sikkim and in Meghalaya (Khasi Hills), and widespread in Nepal and Bhutan; in China only known from western Yunnan (Taylor 1989).

**3.14. *Utricularia subulata* L.:** An annual to perennial terrestrial plant.

**Distribution in northeast India:** Jarain, Meghalaya (Joseph & Mani 1982, misidentified as *U. stanfieldii*, an African species that is not known from India, cf. Taylor 1989).

**Present status:** Widespread and common in India.

**3.15. *Utricularia inflexa* Forssk.:** Medium to large sized suspended aquatic with a whorl of 5-10 floats near the base of the peduncle, corolla white with violet nerves.

**Distribution in northeast India:** Assam (Taylor 1989).

**Present status:** Widespread throughout India.

**3.16. *Utricularia stellaris* L. f.:** Medium to large sized suspended aquatic with a whorl of 3-8 floats above the middle of the peduncle, corolla yellow.

**Distribution in northeast India:** Meghalaya (Joseph & Joseph 1986; Taylor 1989).

**Present status:** Widespread.

**4. *Pinguicula alpina* L. (Lentibulariaceae):** Also known as alpine butterwort, it is a small, herbaceous, perennial insectivore and has white personate spurred flowers, some with a yellow palate.

**Distribution in northeast India:** Sikkim (Singh & Sanjappa 2011).

**Present status:** Widespread in alpine region of the Himalaya.

**5. *Aldrovanda vesiculosa* L. (Droseraceae):** A rare, monotypic, rootless, free floating aquatic carnivorous plant (floats just below the water surface).

**Distribution in northeast India:** Manipur- Imphal (Zaman *et al.* 2011).

**Present status:** Very rare in northeastern India and only once reported from Imphal by Deb in 1965 (Zaman *et al.* 2011) and since then it never have been collected there nor from other parts of India. The IUCN (IUCN 2013) listed the species as Endangered B2ab (iii, v) ver 3.1 (Cross 2012).

**Acknowledgements:** We are thankful to the Director of the Rain Forest Research Institute, Jorhat, Assam, for his encouragement. JS expresses his sincere thanks to Andreas Fleischmann for improving the manuscript.

## References

- Albert, V.A., Williams, S.E., and Chase, M.W. 1992. Carnivorous plants: phylogeny and structural evolution. *Science* 257: 1491-1495.
- Anitha, K. 2011. *Utricularia gibba*. In: IUCN 2013. IUCN Red List of Threatened Species. Version 2013.2.
- Anonymous. 1988. Wealth of India. Vol 3. Council of Scientific and Industrial Research (CSIR), New Delhi.
- Parnell, J.A.N. 2005. An account of the Lentibulariaceae of Thailand. *Thai Forest Bulletin (Botany)* 33: 101-144.
- Basak, R.K. 1975. Distribution of carnivorous plants in West Bengal. *Bull. Bot. Surv. India* 17: 97-107.
- Bhowmik, S., and Datta, B.K. 2012a. Extended distribution of *Utricularia bifida* Linn. (Lentibulariaceae) from India. *Researcher* 4(3): 58-61.
- Bhowmik, S., and Datta, B.K. 2012b. *Utricularia caerulea* L. (Lentibulariaceae) - A new record from Tripura, Northeast India. *NeBIO* 3(1): 35-38.

- Bhowmik, S., and Datta, B.K. 2013. Pollen morphology of some carnivorous plants from Tripura, India. *J. Fundamental Applied Sci.* 2(2): 36-38.
- Bordoloi, R.P.M. 1977. The pitcher plant *Nepenthes khasiana*. Sreeguru Press, Guwahati, India. 59 pp.
- Chew, M.Y., and Haron, N.W. 2011. *Utricularia* (Lentibulariaceae) habitat diversity in Peninsular Malaysia and its implications for conservation. *Gardens' Bulletin Singapore* 63(1 & 2): 451-464.
- Choosawad, D., Leggat, U., Dechusukhum, C., Phongdara, A., and Chotigeat, W. 2005. Anti-tumor activities of fucoidan from the aquatic plant *Utricularia aurea* Lour. *Songklanakarin J. Sci. Technol.* 27(Suppl. 3): 799-807.
- Compton, H. 1909. The morphology and anatomy of *Utricularia brachiata*. *The New Phytologist* 4: 117-130.
- Cross, A. 2012. *Aldrovanda vesiculosa*. In: IUCN 2013. IUCN Red List of Threatened Species. Version 2013.2.
- Deb, D.B. 1983. The Flora of Tripura State, Vols. I & II. Today & Tomorrow's Printers and Publishers, New Delhi.
- Didry, N., Dubreuil, L., Trotin, F., and Pinkas, M. 1998. Antimicrobial activity of aerial parts of *Drosera peltata* Smith on oral bacteria. *Journal of Ethnopharmacology* 60(1): 91-96.
- Dixit, S., and Bera, S.K. 2012. Pollen rain studies in wetland environ of Assam, Northeast India, to interpret present and past vegetation. *International Journal of Earth Sciences and Engineering* 5(4): 739-747.
- Ellison, A.M., and Gotelli, N.J. 2001. Evolutionary ecology of carnivorous plants. *Trends Ecol. Evol.* 16: 623-629.
- Fleischmann, A. 2012. The new *Utricularia* species described since Peter Taylor's monograph. *Carnivorous Plant Newsletter* 41(2): 67-76.
- Gibson, R., Conn, B.J., and Bruhl, J. 2012. Morphological evaluation of the *Drosera peltata* complex (Droseraceae). *Australian Systematic Botany* 25(1): 49-80.
- Hazarika, S., and Barthakur, S.K. 2012. Hydrophytic flora of Assam – I. five new records. *Pleione* 6(2): 401-405.
- IUCN. 2013. IUCN Red List of Threatened Species. Version 2013.2. <http://www.iucnredlist.org>, accessed 12 December 2013.
- Jain, S.K., and Sastry, A.R.K. 1980. Threatened plants of India – a state-of-the-art report. Botanical Survey of India, Howrah.
- Janarthanam, M.K., and Henry, A.N. 1992. Bladderworts of India. Botanical Survey of India, Calcutta.
- Jayaram, K., and Prasad, M.N.V. 2006. *Drosera indica* L. and *D. burmanii* Vahl., medicinally important insectivorous plants in Andhra Pradesh-regional threats and conservation. *Current Science* 91: 943-946.
- Jebb, M.H.P., and Cheek, M.R. 1997. A skeletal revision of *Nepenthes* (Nepenthaceae). *Blumea* 42(1): 1-106.
- Joseph, J., and Joseph, K.M. 1986. Insectivorous plants of Khasi and Jaintia Hills, Meghalaya, India: A preliminary survey. Botanical Survey of India, Calcutta.
- Joseph, J., and Mani, J. 1982. One new species and one new record from India in the genus *Utricularia*. *Bull. Bot. Surv. India* 24(1-4): 108-111.
- Juniper, B.E., Robins, R.J., and Joel, D.M. 1989. The Carnivorous Plants. Academic Press, San Diego, California.
- Kamble, M.V., Harikrishnan, S., and Balakumar, P. 2012. *Utricularia caerulea* (Lentibulariaceae): a



- new record to flora of Andaman & Nicobar Islands. *Rheedea* 22(2): 116-118.
- Kato, Y., Mochida, M., and Koike, H. 2006. The northernmost locality of *Utricularia caerulea* L. (Lentibulariaceae) in Japan. *The Journal of Japanese Botany* 81(1): 41-43.
- Köppen, W., and Geiger, G. (eds.) 1930–1939. *Handbuch der Klimatologie*. 5 vols., Borntraeger, Berlin.
- Lal, J., Chaturvedi, S.K., and Shukla, B.K. 2003. *Utricularia striatula* Smith (Fam. Lentibulariaceae) from Nagaland. Abstract. 5th Indian Agricultural Scientists and Farmers' Congress, 15-16 Feb. 2003, University of Allahabad, Bioed Research and Communication Centre, Allahabad, India.
- Lansdown, R.V., Knees, S.G., and Patzelt, A. 2013. *Utricularia striatula*. In: IUCN 2013. IUCN Red List of Threatened Species. Version 2013.2.
- Li, Z.Y., and Cheek, M.R. 2011. Lentibulariaceae. *Flora of China* 19: 480-491.
- Majumdar, K., Datta, B.K., and Shankar, U. 2011. Community structure and population status of *Drosera burmanii* Vahl. with new distributional record in Tripura, India. *Journal of Ecology and the Natural Environment* 3(13): 410-414.
- Mandal, A., and Mukherjee, A. 2011. *Nepenthes khasiana*: the pitcher plant needs attention for conservation. *Current Science* 100(6): 807.
- Natarajan, K., Kottaimuthu, R., Balasubramanian, V., Pandian, P.A., Malaisamy, M., and Ponnuchamy, A. 2008. Note on the identity of carnivorous plants of Karungalakudi, Tamil Nadu, India. *Ethnobotanical Leaflets* 12: 1073-1077.
- Oliver, D. 1859. III. The Indian species of *Utricularia*. *J. Proc. Linn. Soc., Bot.* 3: 170-190.
- Panday, S., Sinha, B.K., and Karmakar, P. 2013. Three new additions for the state flora of Mizoram, India. *Pleione* 7(1): 266-269.
- Parnell, J.A.N. 2005. An account of the Lentibulariaceae of Thailand. *Thai Forest Bulletin (Botany)* 33: 101-144.
- Prasad, M.N.V., and Jeeva, S. 2009. Coal mining and its leachate are potential threats to *Nepenthes khasiana* Hook. f. (Nepenthaceae) that preys on insects - an endemic plant in North Eastern India. *Biological Diversity and Conservation* 2/3: 29-33.
- Raju A., and Christina, A.J.M. 2013. *Drosera burmannii* Vahl: Antioxidant potential in Dalton's Ascites Lymphoma (DAL) bearing mice. *Journal of Medicinal Plant Studies* 1(4): 152-159.
- Rao, A.S., and Joseph, J. 1967. *Utricularia pubescens* Sm. – first report of its occurrence in India. *Indian Forester* 93: 32-33.
- Rathore, T.S., Tandon, P., and Shekhawat, N.S. 1991. *In vitro* regeneration of pitcher plant (*Nepenthes khasiana* Hook. f.) – a rare insectivorous plant of India. *J. Plant Physiol.* 139: 246-248.
- Rice, B.A. 2006. *Growing Carnivorous Plants*. Timber Press, Portland, USA.
- Santapau, H., and Henry, A.N. 1976. *A dictionary of the flowering plants in India*. Publication and Information Directorate, New Delhi.
- Saxena, H.O. 1965. *Utricularia pubescens* Sm. – a new record for India. *Indian Forester* 91: 73-75.
- Sharma, B.K. 2012. Phytoplankton diversity of a floodplain lake of the Brahmaputra River basin of Assam, north-east India. *Indian J. Fish.* 59(4): 131-139.
- Singh, B., Phukan, S.J., Sinha, B.K., Singh, V.K., and Borthakur, S.K. 2011. Conservation strategies for *Nepenthes khasiana* in the Nokrek Biosphere Reserve of Garo hills, Northeast, India. *International Journal of Conservation Science* 2(1): 55-64.
- Singh, P., and Sanjappa, M. 2011. Flowering plants of Sikkim - An analysis. In: *Biodiversity of Sikkim - Exploring and Conserving a Global Hotspot*. Arrawatia, M.L., and Tambe, S. (eds.). Department of Information and Public Relations, Government of Sikkim, Gangtok. pp. 65-88.

- Subramanyam, K., and Kamble, N. P. 1968. Chromosome numbers in certain Indian species of *Utricularia* L. (Lentibulariaceae). Proceedings of the Indian Academy of Sciences - Section B, 68(5): 221-224.
- Suksathan, P., and Parnell, J.A.N. 2010. Three new species and two new records of *Utricularia* L. (Lentibulariaceae) from Northern Thailand. Thai For. Bull. (Bot.) 38: 23-32.
- Takhtajan, A.L. 1986. Floristic Regions of the World. University of California Press, Berkeley and Los Angeles, California.
- Taylor, P. 1989. The genus *Utricularia* - a taxonomic monograph. Kew Bulletin Additional Series XIV: London.
- Venugopal, N., and Devi, N.R. 2003. Development of the anther in *Nepenthes khasiana* Hook.f. (Nepenthaceae), an endemic and endangered insectivorous plant of North East India. Feddes Repertorium 114(1-2): 69-73.
- Venugopal, N., Devi, R.K., and Rao, C.S. 2007. An interesting observation on the mycorrhizal symbiosis in the insectivorous plant, *Drosera peltata* Sm. in Meghalaya, North-East India. Carnivorous Plant Newsletter 36(1): 9-13.
- Wikipedia. 2013. [http://en.wikipedia.org/wiki/List\\_of\\_Nepenthes\\_species](http://en.wikipedia.org/wiki/List_of_Nepenthes_species), accessed 11 December 2013.
- Yee, C.M., Rahman, R.A., and Haron, N.W. 2012. Micro-structure studies on *Chirita* and *Utricularia* of peninsular Malaysia. Pak. J. Bot. 44(6): 2063-2066.
- Zaman, M., Naderuzzaman, A.T.M., Hasan, M., and Naz, S. 2011. Ecology, morphology and anatomy of *Aldrovanda vesiculosa* L. (Droseraceae) from Bangladesh. Bangladesh J. Bot. 40(1): 85-91.
- Zhuang, X. 2011a. *Drosera peltata*. In: IUCN 2013. IUCN Red List of Threatened Species. Version 2013.2.
- Zhuang, X. 2011b. *Utricularia bifida*. In: IUCN 2013. IUCN Red List of Threatened Species. Version 2013.2
- Ziemer, B. 2010. Exciting conservation news: The Rare *Nepenthes* Collection project! Carnivorous Plant Newsletter 39(3): 67.

## Carnivorous Plant Nursery

*Promoting environmental stewardship through  
engaging educational materials and low-cost, high-quality carnivorous plants.*



- Carnivorous Plants
- Bog Orchids, Bog Plants
- Live Wingless Fruitflies
- Activity Book for Carnivorous Plants
- Bugs in the Bog Board Game
- Free Monthly Drawing
- Online Shopping and Mail Order

**[www.carnivorousplantnursery.com](http://www.carnivorousplantnursery.com)**

301-519-7505 | [info@carnivorousplantnursery.com](mailto:info@carnivorousplantnursery.com) | 16128 Deer Lake Road, Derwood, MD 20855